

Novalac Allergy

Scientifically formulated for babies with Cow's Milk Protein Allergy

Infant formula from birth to 12 months

Novalac Allergy has been scientifically formulated with protein sourced from rice. Novalac Allergy is the first and only 100% rice protein-based infant formula available in Australia and is suitable for babies who are unable to tolerate formula containing cow's milk protein, or with a confirmed or suspected cow's milk protein allergy. Novalac Allergy is also suitable for lactose-intolerant babies. Novalac Allergy nourishes baby with essential daily nutrients and minerals and is suitable for ongoing everyday use. Australasian society of clinical immunology and allergy (ASCIA) lists rice-based formula as an alternative milk for infants (<1yr) with a cow's milk protein allergy.¹

Composition per 100mL Novalac Allergy formula (reconstituted at one 4.5 g scoop / 30mL water)

Average Values	Per 100ml prepared feed*		Average Values	Per 100ml prepared feed*	
Energy	68	kcal	Chromium	<5	µg
	284	kJ	Molybdenum	<7	µg
Protein equivalent	1.8	g	Vitamins		
Carbohydrates	7.4	g	Vitamin A	60.8	µg RE
Lactose	<0.00135	g	Vitamin B1	54	µg
Galactose	<0.00135	g	Vitamin B2	108	µg
Fats	3.4	g	Vitamin B6	40.5	µg
Linoleic Acid [Omega 6]	0.6	g	Vitamin B12	0.2	µg
-Linolenic Acid [Omega 3]	63.3	mg	Vitamin C	8.1	mg
ARA	10.8	mg	Vitamin D3	0.9	µg
DHA	10.5	mg	Vitamin E	1.5	mgα-TE
Minerals			Vitamin K1	6.1	µg
Sodium	27	mg	Niacin	0.6	mg
Potassium	67.5	mg	Pantothenic acid	0.3	mg
Chloride	47.3	mg	Folic acid	8.1	µg
Calcium	60.8	mg	Biotin	2	µg
Phosphorus	33.8	mg	Other Nutrients		
Magnesium	6.1	mg	Total Nucleotide 5' monophosphates	3.2	mg
Iron	0.7	mg	Choline	8.1	mg
Zinc	0.7	mg	Inositol	3.6	mg
Iodine	8.8	µg	Taurine	5.4	mg
Copper	47.3	µg	L-carnitine	1.1	mg
Manganese	13.5	µg			
Selenium	1.5	µg			

*As prepared according to instructions

Novalac Allergy ingredients: Glucose syrup, vegetable oils (palm, rapeseed, sunflower, coconut), rice proteins hydrolysate, corn starch, emulsifier (sunflower lecithin), tricalcium phosphate, monocalcium phosphate, calcium chloride, L-lysine, long chain polyunsaturated fatty acids ARA (single cell oil source) and DHA (fish oil), magnesium sulphate, trisodium citrate, vitamins (A, B1, B12, B2, B6, C, D3, E, K1, biotin, folic acid, niacin, pantothenic acid), L-tryptophan, L-cystine, choline bitartrate, L-isoleucine, ferric pyrophosphate, taurine, nucleotides (cytidine-5'-monophosphate, uridine-5'-monophosphate sodium salt, adenosine-5'-monophosphate, guanosine-5'-monophosphate sodium salt, inosine-5'-monophosphate sodium salt), inositol, zinc sulphate, L-carnitine, antioxidant: extract rich in tocopherols, sodium selenite, cupric sulphate, potassium chloride, potassium iodide, manganese sulphate, calcium citrate, potassium citrate. Contains: Fish

Breast milk is best for babies. Professional advice should be followed before using an infant formula. Introducing partial bottle feeding could negatively affect breastfeeding. Good maternal nutrition is preferred for breastfeeding and reversing a decision not to breastfeed may be difficult. Infant formula should be used as directed. Proper use of an infant formula is important to the health of the infant. Social and financial implications should be considered when selecting a method of feeding.

References: 1. Australasian Society of Clinical Immunology and Allergy. Cow's milk (dairy) allergy. 2019.



SPECIALIST IN INFANT FEEDING CONDITIONS

