

Novalac Colic

Scientifically formulated for babies with colic

Infant formula from birth to 12 months

Novalac Colic is a premium infant formula scientifically formulated to help provide relief to babies with symptoms of colic. Colic is a source of distress for parents, carers and babies alike. For many babies, the bloating and wind associated with colic may be related to undigested lactose that remains in the gut after feeding. Novalac Colic nourishes baby with essential daily nutrients and minerals and is suitable for ongoing everyday use after colic symptoms have been resolved.

Nutritional Information per 100 mL Novalac Colic formula

(reconstituted at one 4.3 g scoop/30mL water)

Average Values	Per 100ml prepared feed*		Average Values	Per 100ml prepared feed*	
Energy	65	kcal	Vitamins		
	274	kJ	Vitamin A	58.5	µg RE
Proteins	1.6	g	Vitamin B1	52	µg
Carbohydrates	7.3	g	Vitamin B2	130	µg
Fats	3.3	g	Vitamin B6	39	µg
Linoleic Acid [Omega 6]	0.5	g	Vitamin B12	0.2	µg
α-Linolenic Acid [Omega 3]	53.3	mg	Vitamin C	7.8	mg
Minerals			Vitamin D3	1	µg
Sodium	21.5	mg	Vitamin E	1.1	mgα-TE
Potassium	68.9	mg	Vitamin K1	3.9	µg
Chloride	46.8	mg	Niacin	0.6	mg
Calcium	65	mg	Pantothenic acid	0.3	mg
Phosphorus	39	mg	Folic acid	7.8	µg
Magnesium	5.9	mg	Biotin	2	µg
Iron	0.9	mg	Other Nutrients		
Zinc	0.5	mg	Choline	10.4	mg
Iodine	10.4	µg	Inositol	5.2	mg
Copper	45.5	µg	Taurine	5.7	mg
Manganese	10.4	µg	L-carnitine	1	mg
Selenium	1.6	µg			

*As prepared according to instructions

Novalac Colic ingredients: Skimmed cow's milk, maltodextrin, vegetable oils (palm, sunflower, soy, rapeseed), milk proteins, lactose, tricalcium phosphate, tricalcium citrate, disodium phosphate, emulsifier (soy lecithin), potassium chloride, vitamins (A, B1, B12, B2, B6, C, D3, E, K1, biotin, folic acid, niacin, pantothenic acid), choline bitartrate, magnesium chloride, dipotassium phosphate, taurine, ferrous sulphate, inositol, L-cystine, magnesium oxide, zinc sulphate, L-tryptophan, antioxidants (extract rich in tocopherols), L-carnitine, potassium odide, sodium selenite, cupric sulphate, manganese sulphate. Contains: Milk, soy.

Breast milk is best for babies. Professional advice should be followed before using an infant formula. Introducing partial bottle feeding could negatively affect breastfeeding. Good maternal nutrition is preferred for breastfeeding and reversing a decision not to breastfeed may be difficult. Infant formula should be used as directed. Proper use of an infant formula is important to the health of the infant. Social and financial implications should be considered when selecting a method of feeding.



SPECIALIST IN INFANT FEEDING CONDITIONS

