

# Novalac Diarrhoea

Scientifically formulated for babies with diarrhoea

## Infant formula from birth to 12 months

Novalac Diarrhoea is a premium infant formula that has been scientifically formulated to provide nutritional support for babies with diarrhoea. Novalac Diarrhoea is low in lactose and contains electrolytes to help rehydrate. Novalac Diarrhoea is for short term use only (up to 5 days exclusively). Once baby's diarrhoea has resolved, gradually re-introduce the baby's regular formula.

### Nutritional Information per 100 mL Novalac Diarrhoea formula (reconstituted at one 4.3 g scoop/30 mL water)

Average Values	Per 100ml prepared feed*	
<b>Energy</b>	<b>62.3</b>	<b>kcal</b>
	<b>261.2</b>	<b>kJ</b>
<b>Protein equivalent</b>	<b>2.3</b>	<b>g</b>
<b>Carbohydrates</b>	<b>7.1</b>	<b>g</b>
Lactose	<0.0013	g
Galactose	<0.0013	g
<b>Fats</b>	<b>2.7</b>	<b>g</b>
Linoleic Acid [Omega 6]	0.5	g
$\alpha$ -Linolenic Acid [Omega 3]	49.4	mg
<b>Minerals</b>		
Sodium	31.2	mg
Potassium	83.2	mg
Chloride	49.4	mg
Calcium	50.7	mg
Phosphorus	26	mg
Magnesium	6.5	mg
Iron	0.1	mg
Zinc	0.5	mg
Iodine	6.5	$\mu$ g
Copper	45.5	$\mu$ g
Manganese	26	$\mu$ g
Selenium	1.3	$\mu$ g

Average Values	Per 100ml prepared feed*	
<b>Vitamins</b>		
Vitamin A	58.5	$\mu$ g RE
Vitamin B1	52	$\mu$ g
Vitamin B2	104	$\mu$ g
Vitamin B6	45.5	$\mu$ g
Vitamin B12	0.2	$\mu$ g
Vitamin C	7.8	mg
Vitamin D3	1	$\mu$ g
Vitamin E	1.4	mg $\alpha$ -TE
Vitamin K1	3.9	$\mu$ g
Niacin	0.6	mg
Pantothenic acid	0.3	mg
Folic acid	7.8	$\mu$ g
Biotin	2	$\mu$ g
<b>Other Nutrients</b>		
Choline	5.2	mg
Inositol	3.3	mg
Taurine	5.7	mg
L-carnitine	1.2	mg

\*As prepared according to instructions

**Novalac Diarrhoea ingredients:** Banana puree, maltodextrins, vegetable oils (palm, soy, coconut), apple puree, milk proteins, partially hydrolysed rice flour, fructose, dextrose, calcium chloride, calcium carbonate, sodium chloride, trisodium citrate, emulsifier (soy lecithin), tricalcium phosphate, vitamins (A, B1, B12, B2, B6, C, D3, E, K1, biotin, folic acid, niacin, pantothenic acid), L-cystine, magnesium sulphate, taurine, disodium phosphate, choline chloride, inositol, L-carnitine, zinc sulphate, antioxidants (extract rich in tocopherols), ferrous sulphate, cupric sulphate, potassium iodide, manganese sulphate, sodium selenite. Contains: Soy, milk.

**Breast milk is best for babies.** Professional advice should be followed before using an infant formula. Introducing partial bottle feeding could negatively affect breastfeeding. Good maternal nutrition is preferred for breastfeeding and reversing a decision not to breastfeed may be difficult. Infant formula should be used as directed. Proper use of an infant formula is important to the health of the infant. Social and financial implications should be considered when selecting a method of feeding.



SPECIALIST IN INFANT FEEDING CONDITIONS

